

- Turner JE, Novacky A (1974) The quantitative relation between plant and bacterial cells involved in the hypersensitive reaction. *Phytopathology* 64: 885-890
- Van Gijsegem F (1986) Analysis of the pectin-degrading enzymes secreted by three strains of *Erwinia chrysanthemi*. *J Gen Microbiol* 132: 617-624
- Van Gijsegem F (1989) Relationship between the *pel* genes of the *pelADE* cluster in *Erwinia chrysanthemi* strain 8374. *Mol Microbiol* 3: 1415-1424
- Van Gijsegem F, Genin S, Boucher C (1993) Evolutionary conservation of pathogenicity determinants among plant and animal pathogenic bacteria. *Trends Microbiol* 1: 175-180
- Verda ME, Scheffler HJ (1988) Hypoxic stress inhibits the appearance of wound-response proteins in potato tubers. *Plant Physiol* 88: 805-809
- Verda ME, Antonov LS, Yang Z, Butler WO, Lacey GH (1992) Hypoxic stress inhibits aerobic wound-induced resistance and activates hypoxic resistance to bacterial soft rot. *Am Potato J* 69: 239-253
- Wandersman C (1989) Secretion, processing and activation of bacterial extracellular proteases. *Mol Microbiol* 3: 1825-1831
- Wandersman C, Delpechere P, Letoffe S, Schwartz M (1987) Characterization of *Erwinia chrysanthemi* extracellular proteases: cloning and expression of the protease genes in *Escherichia coli*. *J Bacteriol* 169: 5046-5053
- Wei ZM, Luby RJ, Zunhoff CH, Bauer DW, Ito SY, Collmer A, Beer SV (1992) Harpin, elicitor of the hypersensitive response produced by the plant pathogen *Erwinia amylovora*. *Science* 257: 85-88
- Went MC, Inness RM, Bent AF, Staskiewicz BJ (1991) Identification of *Pseudomonas syringae* pathogens of Arabidopsis and a bacterial locus determining virulence on both Arabidopsis and soybean. *Plant Cell* 3: 48-59
- Willis DK, Balle TM, Kinscherf (1991a) Genetics of toxin production and resistance in phytopathogenic bacteria. *Experientia* 47: 765-771
- Willis DK, Rich JJ, Hrabak EM (1991b) *hrp* genes of phytopathogenic bacteria. *Mol Plant Microbe Interact* 4: 132-138
- Xiao Y, Lu Y, Heu S, Hutcherson SW (1992) Organization and environmental regulation of the *Pseudomonas syringae* *pv. syringae* 61 *hrp* cluster. *J Bacteriol* 174: 1734-1741
- Yamazaki N, Fry SC, Darvill AG, Albersheim P (1983) Host-pathogen interactions. XXIV. Fragments isolated from suspension-cultured sycamore cell walls inhibit the ability of the cells to incorporate [<sup>14</sup>C] leucine into proteins. *Plant Physiol* 72: 884-889
- Yang Z, Cramer CL, Lacey GH (1992) *Erwinia carotovora* subsp. *carotovora* pectate lyases: In plants gene activation and roles in soft-rot pathogenesis. *Mol Plant Microbe Interact* 5: 104-112
- Yoder MD, Keen NT, Jurtek F (1993) New domain motif: the structure of pectate lyase C, a secreted plant virulence factor. *Science* 260: 1503-1507
- Young JM (1974) Development of bacterial populations in vivo in relation to plant pathogenicity. *NZ J Agric Res* 17: 105-113
- Yucel I, Xiao Y, Hutcherson SW (1989) Influence of *Pseudomonas syringae* culture conditions on initiation of the hypersensitive response of cultured tobacco cells. *Appl Environ Microbiol* 55: 1724-1729
- Zink RI, Engwald JK, McFeyoy JJ, Chatterjee AK (1985) *reca* is required in the induction of pectin lyase and carotolysin in *Erwinia carotovora* subsp. *carotovora*. *J Bacteriol* 164: 390-398
- Zur M, Hankin L (1970) Regulation of pectate lyase synthesis in *Pseudomonas fluorescens* and *Erwinia carotovora*. *J Bacteriol* 104: 13-18
- Zucker M, Hankin L, Sands D (1972) Factors governing pectate lyase synthesis in soft rot and non-soft rot bacteria. *Physiol Plant Pathol* 2: 59-67

## *hrp* genes of Phytopathogenic Bacteria

U. BOWEN

Bowen, U. 1994. *hrp* genes of phytopathogenic bacteria. Pages 79-98 in: Current Topics in Microbiology and Immunology, Vol. 192: Bacterial Pathogenesis of Plants and Animals - Molecular and Cellular Mechanisms. J. L. Dangl, ed. Springer-Verlag, Berlin.

1	Introduction	79
2	Isolation of <i>hrp</i> Genes and General Features	80
3	Structural Organization and Relatedness of <i>hrp</i> Clusters	83
4	Function of <i>hrp</i> Genes in Xanthomonas campestris <i>pv.</i> vesicatoria and Other Plant Pathogenic Bacteria	83
5	<i>hrp</i> -dependent Secretion of Hypersensitive Response-Inducing Proteins	89
5.1	Harpin from <i>Erwinia amylovora</i>	89
5.2	Harpin from <i>Pseudomonas syringae</i> <i>pv. syringae</i>	90
5.3	PopA from <i>Pseudomonas solanacearum</i>	90
6	Regulation of Expression of <i>hrp</i> Genes	91
6.1	<i>Pseudomonas syringae</i>	91
6.2	Regulatory Genes <i>hrpS</i> and <i>hrpN</i> of <i>Pseudomonas syringae</i> <i>pv. phaseolicola</i>	91
6.3	Conserved Sequence Boxes in <i>Pseudomonas syringae</i>	92
6.4	<i>Xanthomonas campestris</i>	93
6.5	<i>Erwinia</i> and <i>Pseudomonas solanacearum</i>	94
	References	94

## 1 Introduction

In nature plants are resistant to the majority of pathogens, and many bacteria live in close contact with the plant without causing any harm (see chapter by Beattie and Lindow in this volume). Among the 1600 different species known in the bacterial kingdom only a small number (about 80) are plant pathogenic and in most cases highly specialized with respect to the plant that can be attacked. Only a few of these species are gram-positive, e.g., *Clavibacter* spp. and *Streptomyces* spp. In this review I focus on subspecies of the gram-negative genera *Erwinia*, *Pseudomonas*, and *Xanthomonas*, which comprise the major bacterial plant pathogens.

To be a successful pathogen the invading bacterium has to overcome the plant's defense. During evolution plant pathogenic bacteria have acquired multiple functions that enable them to colonize and multiply in living plant tissue. In nature, bacteria enter the plant through natural openings (stomata, hydathodes) or

wounds. The bacterial armory contains a number of weapons that contribute to pathogenicity. Obvious examples include degradative extracellular enzymes such as pectinases, cellulases, and proteases. When the corresponding genes are mutated, bacterial ability to invade plant tissues is more or less affected depending on the pathogen, i.e., these functions contribute to and modulate development and severity of infection to different extents (see chapters by Dow and Daniels, and Collmer and Bauer in this volume).

In addition, phytopathogenic bacteria possess a large number of genes needed for basic pathogenicity. These genes have been operationally defined as *hrp* (hypersensitive reaction and pathogenicity; LUNDGREN et al. 1986) based on their mutant phenotype. *hrp* genes are not only essential for pathogenicity on a plant, i.e., the ability to cause disease in a compatible interaction, but also for the incompatible interaction with resistant host varieties or with plants that are not normally a host for the particular pathogen (so called non-host). The incompatible interaction is often associated with the induction of a hypersensitive reaction (HR) in the plant. In contrast to the use of the term hypersensitivity in the animal field, in plants the HR is a rapid defense response involving localized plant cell death, production of phenolics and antimicrobial agents, e.g., phytoalexins, at the site of infection (KLEMER 1982; LUNDGREN et al. 1993). The HR results in prevention of pathogen multiplication and spread and thus in prevention of disease development. Under natural infection conditions the HR is microscopically small and can be induced by just one bacterial cell. Only when bacteria are introduced into plant tissue at high cell densities in the laboratory (about  $10^8$  colony forming units or more/ml) is the HR macroscopically visible as confluent necrosis and can be clearly distinguished from typical disease symptoms. It is important to note that saprophytic or nonpathogenic bacteria such as *Escherichia coli* or *Pseudomonas fluorescens* do not induce the HR and are unable to multiply in plant tissue.

## 2 Isolation of *hrp* Genes and General Features

*hrp* genes have been isolated from all major gram-negative plant pathogenic bacteria except *Agrobacterium*. There are excellent reviews that describe the early work on loci more on one particular pathogen (WILKUS et al. 1991; BOUCHER et al. 1992). The majority of *hrp* genes have been identified by complementation of loss-of-function mutants. Mutants obtained by random chemical (e.g., N-methyl-N'-nitro-N-nitrosoguanidine) or transposon mutagenesis of a pathogenic wild-type strain were inoculated into the host plant and screened for loss of both the ability to cause disease in susceptible plants and to induce the HR in resistant host or non-host plants (often tobacco). The second criterion for the isolation of genes specific for the plant interaction was to ensure that the mutants would still grow in minimal medium. This way mutants affected in genes for basic housekeeping functions were eliminated. A third characteristic of all *hrp* mutants is that they are unable to grow in the plant.

The *hrp* genes were originally described for the bean pathogen *Pseudomonas syringae* pv. *phaseolicola*. LUNDGREN and coworkers (1986) isolated *Tn5*-induced mutants of *P. syringae* pv. *phaseolicola* that had lost both the ability to induce halo-blight disease on bean and the HR in tobacco. Complementation with cosmid clones from a genomic library of the wild-type strain resulted in isolation of a cluster of *hrp* genes localized in a 20 kb DNA region. This was the first indication that both the ability to cause disease and to induce the HR are mediated by common steps in a "pathway".

Since then *hrp* gene clusters have been cloned from a number of different bacteria. Examples include *Pseudomonas solanacearum* (BOUCHER et al. 1987; Fig. 1B), the *Xanthomonas campestris* pathogens *campestris* and *vitiensis* (AHLARI et al. 1991), *translucens* (MAYER et al. 1991), and *vesicatoria* (BOWS et al. 1991; Fig. 1A), *Erwinia amylovora* (STREIBERGER and BEER 1988; BAUER et al. 1990; WALTERS et al. 1990; BAUER and BEER 1991), and several other pathogens of *P. syringae* (e.g., HUANG et al. 1988; LUNDGREN et al. 1988; Fig. 1C). In addition, genes with DNA homology, and in some cases functional homology, have been isolated from other species, e.g., the so-called *wts* genes from *E. stewartii* (CORLIE et al. 1992; LAVER and BEER 1992), and a region containing pathogenicity genes from *X. c. pv. glycines* that complement *hrp* mutants of *X. c. pv. vesicatoria* (HUANG et al. 1992; BOWS, unpublished results). Interestingly, nonpathogenic *xanthomonads* that were originally isolated from diseased plants as opportunists together with pathogenic bacteria do not contain *hrp*-related DNA sequences (SRAU and MUKSAVAGE 1990; BOWS et al. 1991). In *Agrobacterium tumefaciens* or in strains of *Rhizobium* spp. there seem to be no *hrp* gene equivalents present (BOWS et al. 1991; LAVER and BEER 1992). This conclusion is based on DNA hybridization experiments and, of course, does not exclude the presence of genes with functional homology to *hrp* genes in these species.

In all of the cases mentioned above, the *hrp* genes are organized in clusters of 22–40 kb, and I will restrict most of this chapter to these large *hrp* clusters. In addition, several smaller *hrp* loci have been described that are not linked to the large cluster present in the same bacterium. These include a region in *P. solanacearum* (HUANG et al. 1990), the *hrpX* locus that is conserved in *X. campestris* pathogens *campestris* (KAMOUN and KUO 1990; KAMOUN et al. 1992) and *oryzae* (KAMOUN et al. 1993), and the *hrpM* locus in *P. syringae* (NIEPOLO et al. 1985; MUKHOPADHYAY et al. 1988). *hrpM* is functionally conserved in pathovar *phaseolicola* (FELLAY et al. 1991). Besides being nonpathogenic and unable to induce the HR in tobacco, *P. syringae* *hrpM* mutants are also affected in mucus production. The *hrpM* locus encodes two putative proteins which are similar and have been shown to be functionally homologous to the products of the *E. coli* *mdoGH* operon (LOUENS et al. 1993). The *mdoGH* genes are required for periplasmic membrane-derived oligosaccharide synthesis in *E. coli*. The *hrpQ* and *hrpT* genes from *P. syringae* pv. *phaseolicola* (MILLER et al. 1993) will be discussed later in this chapter.




Fig. 1A-C. Genetic and translational organization of the *hrp* gene cluster of different plant-pathogenic bacteria. A *Xanthomonas campestris* pv. *vesicatoria*; B *Pseudomonas solanacearum*; and C *Pseudomonas syringae* pv. *syringae*. Arrows represent transcription units as determined by genetic analyses. Boxes correspond to sequences of open reading frames (ORFs) that have been published. In case of sequence similarities between ORFs in different clusters the boxes are filled with the same pattern. For references, see text.

### 3 Structural Organization and Relatedness of *hrp* Clusters

Genetic studies using transposon-induced insertion mutants in the respective bacterial wild-type strains revealed that the *hrp* clusters contain at least six to eight complementation groups (Fig. 1). Some *hrp* gene clusters have clearly been shown to be localized in the chromosome, e.g., in *P.s. pv. phaseolicola* (Rahme et al. 1991) and in *X.c. pv. vesicatoria* (Bonas et al. 1991), whereas in *P. solanacearum*, the *hrp* cluster is on a megaplasmid (Boucher et al. 1987).

Striking similarities have recently been found between the *hrp* genes of pathogens belonging to different genera. The first indication of homologues came from Southern hybridization studies. DNA homology was observed among different strains of the same pathovar, as well as between pathovars or strains within a species, and in some cases also between species. However, the degree of conservation varies. DNA homology is high within pathovars of a given subspecies, e.g., in *P. syringae* (Lundgren et al. 1988; Huang et al. 1991) and in *X. campestris* (Bonas et al. 1991). The latter studies were recently extended by PCR using primers based on *hrp* sequences from *X.c. pv. vesicatoria* (Lente et al. 1994). Furthermore, at least some regions of the *hrp* clusters appear to be conserved on the DNA level between *P. solanacearum* and pathovars of *X. campestris*, *P. syringae*, and also to *E. amylovora* (Boucher et al. 1987; Arlat et al. 1991; Gough et al. 1992; Laev and Beer 1992). In addition, cross-complementation within a subspecies indicated a high degree of functional conservation of *hrp* genes (e.g., Lindgren et al. 1988; Arlat et al. 1991; Bonas et al. 1991; Laev and Beer 1992). Due to sequence data it is now becoming more and more apparent that several *hrp* genes are conserved in all major grain-negative plant pathogenic bacteria (see below). Whether there are *hrp* genes that are clearly pathovar-specific can only be answered when complete sequence information becomes available for several *hrp* clusters.

### 4 Function of *hrp* Genes in *Xanthomonas campestris* pv. *vesicatoria* and Other Plant Pathogenic Bacteria

DNA sequence analysis of the *hrp* genes has revealed some important clues to their possible biochemical functions. One of the first genes sequenced was a regulatory gene, *hrpS*, from *P.s. pv. phaseolicola* (Grimm and Panopoulos 1989). This gene as well as *hrpB*, a regulatory gene from *P. solanacearum* (Gemm et al. 1992), will be discussed below in the context of gene regulation.

Since *hrp* genes are environmentally regulated (see below), it was believed for a while that they would be encoding "alternative" proteins required for adaptation of the bacterium to the plant as the preferred environment. The recently discovered sequence similarities between several putative Hrp proteins and known proteins from other bacteria, however, led to a very different hypothesis, namely, involvement of Hrp proteins in protein secretion. We have

sequenced the entire *hnp* cluster of *X.c. pv. vesicatoria*. Since most *hnp* sequences from this and other bacteria are not yet published, I will summarize our results and refer to the other phytopathogenic bacteria as I go along. Based on genetic analyses and the open reading frames (ORFs) with a high coding probability we predict 21 *hnp* genes in the 25 kb *hnp* cluster of *X.c. pv. vesicatoria*. Their transcriptional organization is depicted in Fig. 1A. The loci *hnpA* and *hnpB* are transcribed from right to left; the other four loci are transcribed from left to right (Schulte and Bonas 1992a). According to the locus (*hnpA-hnpF*) we have numbered the ORFs consecutively. The *hnpA* locus appears to contain just one *hnp* gene, *hnpA1*. The *hnpB* operon contains eight ORFs, called *hnpB1-hnpB8*, etc. A region of about 4 kb between *hnpE* and *hnpF* does not seem to be involved in the interaction with the plant because insertions in this region do not lead to a change in phenotype (Bonas et al. 1991).

What are the characteristics of the Hrp proteins? It should be noted that, except for three proteins, expression of the other 18 has yet to be demonstrated in *X.c. pv. vesicatoria*. A number of putative Hrp proteins are most likely associated with or localized in the bacterial membrane. For example, the HrpC2 protein sequence contains eight transmembrane domains but lacks a signal sequence, suggesting an inner membrane localization (Fenselau et al. 1992). Both HrpA1 and HrpB3 contain an NH<sub>2</sub>-terminal signal sequence and one (HrpA1) or two (HrpB3) transmembrane domains, suggesting that a part of these proteins might be targeted to the outer membrane. The signal sequence of HrpB3 resembles signal peptidase II sequences which are typical of lipoproteins (Fenselau et al. 1992). Experiments using radioactively labeled palmitate are underway to test whether HrpB3 is a lipoprotein. Recently, both HrpB3 and HrpA1 were shown to be localized in the *X.c. pv. vesicatoria* membrane fraction using polyclonal antibodies (S. Fenselau, C. Marie, and U. Bonas, manuscript in preparation). The HrpB6 protein is a putative ATPase with highly conserved nucleotide and magnesium binding domains. It is more similar to protein traffic ATPases than to proton pump ATPases, and the lack of membrane spanning domains suggests a cytoplasmic location (Fenselau et al. 1992).

Searches of the database revealed sequence relatedness of more than half of the *X.c. pv. vesicatoria* Hrp proteins with putative proteins in other bacteria, including different plant pathogens. High DNA sequence identity (more than 90%) was found to a 2.7 kb fragment carrying pathogenicity genes from *X.c. pv. glycines* (Hwang et al. 1992). The authors predicted two ORFs, whereas in *X.c. pv. vesicatoria*, this region contains three ORFs corresponding to the *hnpC3*, *hnpD1* and *hnpD2* genes. Complementation studies indicated that part of the *hnp* region is colinear in the two pathogens of *Xanthomonas* (unpublished).

The deduced amino acid sequences of *hnp* genes published from *P. solanacearum* (Gough et al. 1992, 1993; Goux et al. 1992) show significant similarity to *X.c. pv. vesicatoria* proteins (Table 1; Fig. 1). One exception is the *hnpB* regulatory gene from *P. solanacearum* which is not present in the 25 kb *hnp* region or in the flanking region of the *X.c. pv. vesicatoria hnp* cluster as determined by DNA sequence analysis and hybridization studies (T. Horis and U. Bonas,

unpublished). Furthermore, several of the proteins mentioned are conserved in other species (Fig. 1), however, the degree of sequence similarity varies greatly (Table 1). The HrpA1 protein from *X.c. pv. vesicatoria* is 48% and 29% identical to proteins from *P. solanacearum* (HrpA; Gough et al. 1992) and *P.s. pv. syringae* (HrpH; Huang et al. 1992), respectively. HrpC2 from *X.c. pv. vesicatoria* is even more conserved, being 66% identical to the corresponding HrpO protein of *P. solanacearum* (Gough et al. 1993), whereas the *hnpI* genes from *E. amylovora* (Wei and Beer 1994) and from *P.s. pv. syringae* (Huang et al. 1993) both show 62% similarity to *hnpC2* from *X.c. pv. vesicatoria*. *P.s. pv. syringae* also contains a *hnpB3* related gene, called *hnpY*, and a *hnpD2* related gene, *hnpW* (H.-C. Huang, personal communication). Thus, the high degree of DNA sequence conservation that was reported earlier (see above) is also seen on the protein level. It appears that *hnp* genes in *X.c. pv. vesicatoria* are more closely related to *P. solanacearum* than to *P. syringae* and to *Erwinia*. As more and more homologous *hnp* genes are found in other bacteria nomenclature might become confusing. However, as long as the genes have not been shown to be functionally homologous, we will continue to use these names.

Besides genes that are conserved among the major phytopathogenic bacteria some genes are absent in the *hnp* region of more distantly related species. For example, there are no known homologs of the harpin genes *hnpN* (Wei et al. 1992a) and *hnpZ* (He et al. 1993) (see below), and of *hnpJ* from *P.s. pv. syringae* (Huang et al. 1993) in the *X.c. pv. vesicatoria hnp* cluster (unpublished; see Fig. 1). Similarities of 50%–60% were found recently between HrpA1 and HrpB3 from *X.c. pv. vesicatoria* and two putative Nol proteins of *Rhizobium fredii* that are encoded by a cultivar specificity region. NolT and NolW mutants have a wider host range in nodulation of soybean (Meuwinkel et al. 1993). In addition, the authors mention that release of proteins is affected.

Last but not least, Table 1 summarizes the significant sequence similarities which have been found to proteins from animal bacterial pathogens. A number of putative Hrp proteins are related to proteins in animal pathogens such as *Salmonella*, *Shigella*, and *Yersinia* spp. Since the first similarities found were to the Ysc, Vir, and Lcr proteins from *Yersinia* spp., this group of organisms became a "role model" for plant pathologists (Fenselau et al. 1992; Gough et al. 1992; Huang et al. 1992). In *Yersinia*, these proteins are essential for the secretion of virulence factors, called Yops (Yersinia outer protein; Michiels et al. 1990, 1991). Since they are described in detail in the chapter by G.-R. Cornelis, I will mention only a few important features. The Yops are hydrophilic proteins that lack a typical NH<sub>2</sub>-terminal signal peptide, and are secreted by using an entirely different pathway from that previously described for protein secretion. The genes involved in secretion are clustered on a 70 kb virulence plasmid. In case of a mutation, e.g., in Ysc, the Yops accumulate in the cytoplasm (McGuire et al. 1991). Although their direct role in transport has yet to be demonstrated, it is believed that the Ysc and Lcr proteins mentioned in Table 1 are parts of a special transport apparatus for Yop secretion. Similarly, *Shigella flexneri* secretes virulence factors, called Ipa (invasion plasmid antigens), that are distinct from Yops but share the general

Table 1. Sequence similarities of *Xanthomonas campestris* pv. *vesicatoria* Hrp proteins

<i>Xanthomonas campestris</i> pv. <i>vesicatoria</i>	HrpA1 <sup>1</sup>	HrpB6 <sup>1</sup>	HrpB3 <sup>1</sup>	HrpC1 <sup>1</sup>	HrpC2 <sup>1</sup>	HrpC3 <sup>1</sup>	HrpD1 <sup>2</sup>	HrpD2 <sup>2</sup>
<i>Pseudomonas solanacearum</i>	HrpA <sup>3</sup> (66%)	HrpE <sup>3</sup>	HrpI <sup>3</sup> (70%)	HrpH <sup>3</sup> (74%)	HrpO <sup>1</sup> (81%)	HrpP <sup>3</sup> (54%)	HrpQ <sup>4</sup>	HrpT <sup>4</sup>
<i>Pseudomonas syringae</i> pv. <i>syringae</i>	HrpH <sup>4</sup> (52%)				HrpI <sup>7</sup> (62%)			
<i>Yersinia enterocolitica</i>	YscC <sup>8</sup> (55%)		YscJ <sup>8</sup> (56%)					
<i>Yersinia pestis</i>	YscC <sup>8</sup> (55%)				LcrD <sup>10</sup> (70%)		LsaA <sup>11</sup> (52%)	LsaB <sup>11</sup> (72%)
<i>Yersinia pseudotuberculosis</i>		YscN <sup>12</sup> (73%)	LcrKa <sup>13</sup> (56%)					
<i>Shigella flexneri</i>	MxdD <sup>14</sup> (50%)	Spa47 <sup>15</sup> (65%)	MxdJ <sup>16</sup> (52%)	Spa40 <sup>17</sup> (55%)	MxdA <sup>18</sup> (65%)			Spa24 <sup>19</sup> (67%)
<i>Salmonella typhimurium</i>	InvG <sup>19</sup> (52%)	SpaL <sup>20</sup> (70%)		SpaS <sup>20</sup> (56%)	InvA <sup>21</sup> (67%)			SpaP <sup>22</sup> (64%)
		FliH <sup>23</sup> (65%)						
<i>Bacillus subtilis</i>		FlaA-ORF4 <sup>24</sup> (68%)		FliH <sup>25</sup> (62%)	FliH <sup>25</sup> (63%)			FliP <sup>26</sup> (68%)
<i>Escherichia coli</i>		β-F1 <sup>27</sup> (53%)						FliP <sup>26</sup> (65%)
<i>Erwinia carotovora</i>							MopB <sup>28</sup> (49%)	MopC <sup>28</sup> (65%)
<i>Erwinia amylovora</i>					HrpI <sup>30</sup> (62%)			
<i>Rhizobium fredii</i>	NolW <sup>31</sup> (51%)		NolT <sup>32</sup> (61%)					
<i>Caulobacter crescentus</i>					FliF <sup>33</sup> (55%)			

Similarities between deduced amino acid sequences of Hrp proteins from *X.c.pv.vesicatoria* and other proteins include conservative amino acid exchanges. Number in parentheses indicates percent similarity.

Superscript numbers indicate references as follows:

1, FENSELAU et al. 1992; 2, Bonas et al., unpublished; 3, GOUGH et al. 1992; 4, GENIN et al. 1993, sequences unpublished; 5, GOUGH et al. 1993; 6, HUANG et al. 1992; 7, HUANG et al. 1993; 8, MICHIELS et al. 1991; 9, HADDIX and STRALEY 1992; 10, PLANO et al. 1991; 11, Fields et al. unpublished, accession # L22495; 12, Galyov, unpublished, accession # U00998; 13, RIMPIAINEN et al. 1992; 14, ALLAOUI et al. 1993; 15, VENKATESAN et al. 1992; 16, ALLAOUI et al. 1992; 17, SASAKAWA et al. 1993; 18, ANDREWS and MAURELU 1992; 19, Lodge et al., unpublished, accession # X75302; 20, GROISMAN and OCHMAN 1993; 21, VOGLER et al. 1991; 22, GALÁN et al. 1992; 23, ALBERTINI et al. 1991; 24, Carpenter et al., unpublished, accession # X741212; 25, CARPENTER and ORDAL 1993; 26, BISCHOFF et al. 1992; 27, SARASTE et al. 1981; 28, MALAKOOTI et al. unpublished, accession # L21994; 29, MULHOLLAND et al. 1993; 30, WEI and BEER 1993; 31, MEINHARDT et al. 1993; 32, RAMAKRISHNAN et al. 1991; SANDERS et al. 1992.




Fig. 2. Hypothetical model of cell signaling between gram-negative bacteria and plants indicating the proposed function of Hrp proteins as an apparatus for protein secretion. The model has been modified after FENSELAV *et al.* (1992). Hrp proteins may form a tunnel that enables the export of molecules such as virulence factors or avirulence factors leading to either a hypersensitive response (HR) or disease. These factors could be encoded by *hrp* genes or genes unlinked to the large cluster. Both types of genes have been found to encode elicitors of the HR (see text). The secretion of virulence proteins is hypothetical.

features mentioned above (Hale 1991; and see chapter by PARSY, this volume). Although *S. typhimurium* appears to possess a secretion system similar to that in *Shigella*, secreted invasion antigens have not yet been identified (GIRJESWARI and OCHIAIM 1993; see chapter by FIKSAVI). As unpublished reports indicate that more and more genes in the animal pathogens are conserved, the data shown in Table 1 will soon be of use. Proteins from other bacteria, e.g., *E. coli*, *Bacillus*, *Caulobacter* and from the *nop* region in *E. carotovora* (MURILLI *et al.* 1993), have also been found to be similar to Hrp proteins (Table 1). Most of these are important for the assembly of the flagella, motility, or chemotaxis, again pointing, in my opinion, to a specialized secretion system rather than an involvement of *hrp* genes in chemotaxis.

These observations led us and others to propose a *hrp*-dependent secretion system in plant pathogenic bacteria (FENSELAV *et al.* 1992; GOUCH *et al.* 1992; VAN GUSST *et al.* 1993). A model is shown in Fig. 2 and raises certain questions, e.g., if secretion occurs, what is being secreted by plant pathogenic bacteria? So far, a few proteins have been identified as elicitors of the HR but there is no evidence for secretion of virulence factors (see below).

## 5 *hrp*-dependent Secretion of Hypersensitive Response-Inducing Proteins

### 5.1 Harpin from *Erwinia amylovora*

An important feature of the isolated *hrp* clusters from both *E. amylovora* and *P. syringae* is the ability of *E. coli* or *Pseudomonas fluorescens* transformants containing the cloned genes to induce the HR on tobacco (HUANG *et al.* 1988; BEER *et al.* 1991; see below). This has prompted to search for the HR-inducing activity within the respective gene clusters.

The first bacterial HR-inducing protein identified, designated harpin, is a cell envelope-associated protein encoded by the *hrpN* gene of *E. amylovora*, a pathogen of pear and apple (WEI *et al.* 1992a). This harpin<sub>N</sub> is a glycine-rich and heat-stable protein that induces the HR in the non-host tobacco. The *hrpN* gene is localized within the respective *hrp* cluster and thus has a dual role in also being required for pathogenicity on the normal host plant. Its function in pathogenicity, however, is unknown. BEER *et al.* (1993) mentioned in a preliminary report that the *hrpN* gene seems to be conserved among *Erwinia* spp. but that there is no DNA homology between *hrpN* and sequences in the other plant pathogenic bacteria. Although data described below suggest that the harpin<sub>N</sub> protein might be secreted via the Hrp secretory apparatus, there is no published information available that demonstrates this.

### 5.2 Harpin from *Pseudomonas syringae* pv. *syringae*

Using an elegant approach He and coworkers recently have identified harpin<sub>PS</sub>, which is encoded by the *hrpZ* gene in the bean pathogen *P. syringae* (He *et al.* 1993; see Fig. 1C and chapter by COLLIER and BAUER). Lysates of *E. coli* clones containing an expression library, made using the cloned *P. syringae* *hrp* cluster, were directly screened for HR-inducing activity on tobacco leaves. Two proteins were identified, one of which was an NH<sub>2</sub>-terminal deletion of harpin<sub>PS</sub> with even higher activity than the full size protein. Whether or not processing occurs in natural infection is not clear. Interestingly, two short direct repeats in the COOH-terminus of harpin<sub>PS</sub> are essential for elicitor activity. Although the two harpins harpin<sub>N</sub> and harpin<sub>PS</sub> differ in their primary sequence, they have several features in common, e.g., a stretch of 22 amino acids that is similar in both proteins (He *et al.* 1993). Harpin<sub>PS</sub> is also glycine-rich and heat-stable. As with harpin<sub>N</sub> of *E. amylovora*, the function of harpin<sub>PS</sub> in pathogenicity is unknown. Its product is secreted by *P. syringae* in a HrpH-dependent way. HrpH is highly related to proteins involved in secretion in other plant and animal pathogens (HUANG *et al.* 1992; see Table 1).

### 5.3 *PopA* from *Pseudomonas solanacearum*

An HR-inducing protein has been identified and characterized from *P. solanacearum* culture supernatants, called Pop (Pseudomonas out protein; Arlat et al. 1994). PopA1 and two shorter derivatives, PopA2 and PopA3, induce the HR in tobacco and in certain, but not all, *Petunia* lines. Like the harpins, the Pop proteins are also heat-stable and glycine-rich, however, the sequence is entirely different. In contrast to the harpins, the *popA* gene is not a *hrp* gene but is located outside of the large *hrp* cluster. Interestingly, expression of *popA* is *hrpB*-dependent, i.e., the gene is part of the *hrp* regulon. Mutations in *popA* do not affect the HR on tobacco or pathogenicity on tomato suggesting that more than one HR-inducing factor is produced. Arlat et al. (1994) convincingly showed that secretion of PopA is dependent on other *hrp* genes, such as *hrpA*, *hrpN*, and *hrpO* (Fig. 1B). If a bacterial strain virulent towards *Petunia* is found it will be interesting to see if PopA acts as an avirulence protein in *Petunia* as has been suggested by the authors.

These exciting findings prove that certain Hrp proteins of *P. syringae* and *P. solanacearum* play a role in transport of HR elicitors (Fig. 2). They also stimulate more questions. It needs to be shown that harpins and PopA are in fact secreted when the bacteria interact with the plant (the *hrp* genes were induced in vitro). Are harpins conserved among pathogens of *P. syringae*? How many elicitors (the *n*-host HR in tobacco can be found)? Is the mechanism of recognition in tobacco identical with the *Erwinia* and *P. syringae* harpins and the *P. solanacearum* Pops?

## 6 Regulation of Expression of *hrp* Genes

Expression of *hrp* genes is controlled by environmental conditions and has been studied on the RNA level as well as using transcriptional fusions to reporter genes, such as the *E. coli* genes encoding  $\beta$ -galactosidase or  $\beta$ -glucuronidase. In general, expression of *hrp* loci is not detectable when the bacteria are grown in complex culture media. However, after growth of the bacteria in the plant, *hrp* genes are expressed. Attempts to mimic the conditions that the different bacterial species encounter in the plant tissue resulted in the finding that growth in minimal media without any plant-derived factor was sufficient to induce *hrp* genes. This has led to the speculation that the bacteria have to experience some kind of starvation conditions for full expression of *hrp* genes. One of the first indications for *hrp* gene expression *in vitro*, and clearly a breakthrough, was a report on the *hrp*-dependent expression of an avirulence gene from the soybean pathogen *P. syringae* (Hurni et al. 1989).

Since the composition of minimal media differs depending on the bacterium studied, the most important findings will be summarized for representative pathogens. Parameters like carbon source, concentration of organic nitrogen and phosphate, osmolarity, and pH have been found to be important. High con-

centration of organic nitrogen generally appears to suppress *hrp* gene activation. Only two regulatory genes have been studied so far (see below). Interestingly, they both belong to different families of regulatory proteins.

### 6.1 *Pseudomonas syringae*

Expression of all seven *hrp* loci in the large cluster of *P. syringae* *phaseolicola* is suppressed in complex medium but induced in the plant. Induction occurs in the susceptible host plant as well as in the non-host, tobacco, suggesting that there is no plant species-specific molecule involved in control of host range (Rahme et al. 1992). Five complementation groups, *hrpAB*, *hrpC*, *hrpD*, *hrpE* and *hrpF*, can also be induced in M9 minimal medium containing sucrose as a carbon source, however, induction is affected by pH, osmolarity, and carbon source, and never reaches the levels obtained in the plant (Rahme et al. 1992). A similar observation was made earlier for the avirulence gene *avrB* in *P. syringae* *glyciniae*. Induction occurred in a minimal medium containing fructose, mannitol, or sucrose. Expression of *avrB* is dependent on *hrp* genes homologous to *hrpRS* and *hrpL* from *P. syringae* *phaseolicola* and was suppressed by TCA cycle intermediates such as citrate and succinate (Hurni et al. 1989). *hrp* gene expression in *P. syringae* occurs in the same medium as described by Hurni et al. (1989). (Huang et al. 1991; Xiao et al. 1992). The authors showed *hrp* gene induction in the non-host plant, tobacco, but no data for the host plant. The *P. syringae* *phaseolicola* loci *hrpL* and *hrpRS* are only expressed to a very low level in M9 minimal medium and are induced at least 1000-fold when the bacteria are inoculated into the plant. This led to the conclusion that, at least for expression of *hrpL* and *hrpRS*, specific plant factors might be necessary (Rahme et al. 1992).

### 6.2 Regulatory Genes *hrpRS* and *rpouN* of *Pseudomonas syringae* pv. *phaseolicola*

The results on environmental factors inducing or suppressing *hrp* gene expression suggested that specific regulatory genes are involved in the control of *hrp* promoter activities. At least two loci are involved in positive regulation of the other *hrp* loci of *P. syringae* *phaseolicola* *hrp* cluster (Fellay et al. 1991). While there is no information published for *hrpL*, *hrpRS* has been sequenced. It contains two genes whose predicted protein products are 60% identical to each other (Grimm and Panopoulos 1989; Miller et al. 1993). The HrpS protein is similar to members of the NtrC family of regulatory proteins in gram-negative bacteria. Most NtrC-like regulatory proteins are members of two-component systems, with a sensor protein that in turn activates a response element by phosphorylation of a site in the conserved NH<sub>2</sub>-terminal domain (Alenxart et al. 1989). The putative sensor component operating in *hrp* gene regulation has not been identified. It is postulated that HrpS is the activating protein, however, direct biochemical data

have not been presented. The lack of a typical NH<sub>2</sub>-terminal domain in HrpS could indicate that a different mechanism may be involved in HrpS activation. Apparently, *hrpS*-related sequences are also present in other bacteria, e.g., in *P. syringae* (Heu and Hureleson 1993) and in *Erwinia amylovora* (Beer et al. 1993). *E. stewartii* contains a transcriptional regulator, *WlsA*, with 52% identity to HrpS of *P. sy. phaseolicola*. The *hrpS* clone, however, was unable to functionally complement a *wlsA* mutant (Frenneick et al. 1993).

The structure of the *hrpRS* locus and the finding of -24/-12 consensus sequences upstream of *hrpRS* indicated a possible role in transcriptional activation for transcription factor sigma 54, encoded by *spoV/GamM* and *Pakopoulos* (1989). In a preliminary report, Feller et al. (1991) demonstrated that *hrp* gene expression in *P. sy. phaseolicola* is indeed dependent on *spoV*. A *spoV* mutant of *P. sy. phaseolicola* is a glutamine auxotroph and nonpathogenic. Whether *spoV* is generally involved in regulation of *hrp* gene expression is not clear. In *X. c. pv. vesicatoria*, *spoV* is clearly not involved in *hrp* gene expression and pathogenicity (T. Horns and U. Bonas, manuscript in preparation).

Recently, Miller et al. (1993) have reported the identification of two new loci, *hrpQ* and *hrpT*, from *P. sy. phaseolicola* that affect activation of *hrpRS* in *trans*. However, since *hrpRS* is strongly induced in plants while both *hrpQ* and *hrpT* are constitutively expressed, there must be more factors involved in *hrp* gene regulation. Strains carrying mutations in either *hrpQ* or *hrpT* are amino acid auxotrophs (methionine and tryptophan). *hrpQ* and *hrpT* are probably involved in methionine and tryptophan biosynthesis, respectively (Miller et al. 1993). As stated above, such mutants would normally have been eliminated from the *hrp* mutant analysis.

### 6.3 Conserved Sequence Boxes in *Pseudomonas syringae*

A conserved sequence, the so-called harp box (TGAC/AANC, Fittler et al. 1991), upstream of four *hrp* loci in *P. sy. phaseolicola*, was suggested to be involved in positive regulation of expression. Similar motifs were described for the promoter regions of several *P. syringae* avirulence genes, the expression of which is dependent on *hrpRS* and on *spoV* (Huyun et al. 1989; Salmeron and Szaszwicz 1993; Innes et al. 1993; Shen and Keen 1993). These studies led to a revised harp box sequence (GGAACCA). Its significance in protein binding has not been shown but *avd* promoter constructs lacking the harp box are no longer inducible (Sien and Keen 1993). A harp box-related motif was also found upstream of transcription unit 3 in *P. solanacearum* (Goun et al. 1993).

There is no harp box sequence in *Xanthomonas* *hrp* gene promoters. Another sequence motif that occurs in the promoter region of *hrp* loci in *X. c. pv. vesicatoria* was recently identified. This "PIP" (plant-inducible promoter) box has the sequence TTCGC-N15-TTCGC and occurs upstream of the -35 consensus sequence in four out of six *hrp* promoters (S. Fenselau and U. Bonas, unpublished). Experiments are underway to test whether this is a protein binding motif.

### 6.4 *Xanthomonas campestris*

Expression of *hrp* genes in *X. c. pv. campestris* was determined after growth in vitro and found to be induced in a minimal medium with sucrose and/or fructose as carbon source. No expression occurred in complex media or with high concentrations of organic nitrogen (Ahar et al. 1991). In *X. c. pv. vesicatoria*, expression of the six *hrp* loci is induced in the plant but cannot be efficiently induced in the synthetic media tested so far. However, culture filtrates of sterile tomato cell suspension cultures (called TCM) induced expression of the six *hrp* loci in *X. c. pv. vesicatoria* whereas the basal Murashige-Skoog culture medium (Schulte and Bonas 1992a). De novo transcription of all *hrp* loci occurs rapidly (within 1 h after transfer of the bacteria into TCM (S. Fenselau and U. Bonas, unpublished). A minimal medium was designed which would not suppress *hrp* gene induction. This medium, called XVM1, induces the *hrpF* locus (Fig. 1A) to high levels and differs from the other media described above, in particular by its low concentration in phosphate. Both sucrose and methionine are needed for efficient induction. It is not known whether a plant factor is necessary for activation of the other *hrp* loci, or if the XVM1 medium still lacks components or contains them in suppressing amounts (Schulte and Bonas 1992b).

### 6.5 *Erwinia* and *Pseudomonas solanacearum*

The *hrp* genes of *Erwinia amylovora* are rapidly induced in the non-host, tobacco, and more slowly in the host, pear. Several loci were induced in minimal medium with mannitol as a carbon source. Induction was suppressed by high concentrations of nitrogen and by glucose and was slightly temperature dependent (Wei et al. 1992b).

In *P. solanacearum*, the *hrp* cluster was also induced in host and in non-host plants, as well as in minimal medium. The best carbon sources for induction of four of the six transcription units were pyruvate and glutamate while, as in other bacteria, casamino acids suppressed induction (Ahar et al. 1992). The two rightmost *hrp* transcription units (5 and 6; Fig. 1B) are constitutively expressed but can be induced under certain conditions (Gemin et al. 1992).

The only other gene reported to regulate *hrp* gene expression is *hrpB* from *P. solanacearum*. The gene is part of the *hrp* cluster and appears to be a member of the AraC family of positive regulatory proteins. Interestingly, *hrpB* is related to *Yersinia* (Comkiet et al. 1989; Gemin et al. 1992). The *hrpB* gene positively regulates four of the six *hrp* loci, as well as the *popA* locus, located outside of the *hrp* cluster which encodes a protein secreted in a Hrp-dependent way (see above; Ahar et al. 1994). Whether the HrpB protein binds directly to *hrp* promoters is not yet known.

At this time one can only speculate whether the regulatory systems for *hrp* gene expression employed by *P. solanacearum* and *P. syringae* are really different



or whether there is a global regulatory network thus allowing the fine tuning of gene expression in response to environmental cues. In conclusion, most *htr* loci from different bacteria are inducible in a particular minimal medium. At this time it cannot be ruled out that stimulation of *htr* gene expression involves specific plant factors as was described for the virulence genes of *Agrobacterium tumefaciens* 1992. Since the composition of the nutrients available to the pathogen in the plant is not known one can only speculate that the conditions described above reflect the situation in the plant. It is noteworthy that the *in vitro* culture will only mimic the dynamic nutritional situation that bacteria experience in their interaction with a plant for a short time. In mammalian bacterial pathogens, the expression of genes involved in virulence is also regulated in response to environmental cues rather than to specific host molecules. This subject has been reviewed recently (Mekalanos 1992 and in accompanying chapters), and I will only mention some important factors. In *Yersinia*, the *vir* and *cr* genes are regulated by low calcium (low calcium response genes, Straley et al. 1993) and by temperature (Cornelius et al. 1989; see chapter by Cornelius). A calcium effect has not been described for any plant bacterium. In our laboratory no effect of calcium on *htr* gene expression in XVM1 was observed (Schulte and U. Bonas, unpublished). Expression of *invA* of *S. typhimurium* of the *mxi* and *ipa* genes of *Shigella* is affected by osmolarity and the later genes also by temperature (Galán and Curtiss 1990; Hale 1991).

**Acknowledgements.** I thank my previous and present coworkers—Ilse Batto, Martina Gutschow, Achille Fenselau, Tosián Horns, Corinne Marie, Michèle Perré, Ralf Schulte, and Kai Wengert—for contributing to the data described here and for fruitful discussions. I am also grateful to my colleagues for sending preprints and sharing unpublished results, and to Heather McKhann and John Mansfield for helpful suggestions on the manuscript. The research in my laboratory was supported in part by grants from the Bundesministerium für Forschung und Technologie (322-4003-0216300A), the Deutsche Forschungsgemeinschaft, and the EEC (BIO1-CT90-0168).

## References

- Albertini AM, Caramori I, Crabo WD, Scollone F, Gahrz A (1991) The *lba* locus of *Bacillus subtilis* is part of a large operon coding for flagellar structures, motility functions, and an ATPase-like polypeptide. *J Bacteriol* 173: 3573-3579
- Albright LM, Huala E, Ausubel FM (1989) Prokaryotic signal transduction mediated by sensor and regulator protein pairs. *Annu Rev Genet* 23: 311-336
- Allouit A, Sansonetti PJ, Paros C (1992) MutA, a lipoprotein involved in secretion of *Shigella flexneri*, is homologous to YnfJ, a structural factor of the *Yersinia* Yop proteins. *J Bacteriol* 174: 7761-7769
- Altieri A, Sansonetti PJ, Paros C (1993) MutA, an outer membrane protein necessary for the secretion of the *Shigella flexneri* *ipa* invasins. *Mol Microbiol* 7: 59-68
- Autreau GP, Mauriell AT (1992) *mutA* of *Shigella flexneri* 2a, which facilitates export of invasion plasmid antigens, encodes a homolog of the low-calcium response protein, *LeuO*, of *Yersinia* *pasteurii*. *Infect Immun* 60: 3287-3295
- Ayali M, Gough CL, Barber CE, Boucher C, Daniels MJ (1991) *Xanthomonas campestris* contains a cluster of *htr* genes related to the *htr* cluster of *Pseudomonas solanacearum*. *Mol Plant Microbe Interact* 4: 593-601
- Ayali M, Gough CL, Zischek C, Barberis PA, Tigelet A, Boucher CA (1992) Transcriptional organization and expression of the large *htr* gene cluster of *Pseudomonas solanacearum*. *Mol Plant Microbe Interact* 5: 187-193

- Ayali M, Van Gijsegem F, Hual JC, Pernodet JC, Boucher CA (1994) *PopA1*, a protein which induces a hypersensitive-like response on specific *Petunia* genotypes, is secreted via the *Hrp* pathway of *Pseudomonas solanacearum*. *EMBO J* 13: 543-553
- Barry MA, Guiberville MH, Marcis B, Colasse A, Paulin JP, Laurent J (1990) Cloning of a large gene cluster involved in *Erwinia amylovora* CFBP1430 virulence. *Mol Microbiol* 4: 777-787
- Bauer DW, Beer SV (1991) Further characterization of an *htr* gene cluster of *Erwinia amylovora*. *Mol Plant Microbe Interact* 4: 483-489
- Beer SV, Bauer DW, Jiang X, Luby RJ, Sneath BJ, Weil ZM, Wilcox DA, Zurnoff CH (1991) The *htr* gene cluster of *Erwinia amylovora*. In: Hamecke H, Verma DPS (eds) *Advances in molecular genetics of plant-microbe interactions*, vol 1, Kluwer Academic, Dordrecht, Netherlands, pp 53-60
- Beer SV, Weil ZM, Luby RJ, He SY, Bauer DW, Colmer A, Zurnoff CH (1993) Are *htr* genes universal elicitors of the hypersensitive response of phytopathogenic bacteria? In: Nestler EW, Verma DPS (eds) *Advances in molecular genetics of plant-microbe interactions* vol 2, Kluwer Academic, Dordrecht, Netherlands, pp 281-286
- Beschall DS, Weinreich MR, Odel GW (1992) Nucleotide sequence of *Bacillus subtilis* flagellar biosynthetic genes *flp* and *flhD* and identification of a novel flagellar gene. *J Bacteriol* 174: 4017-4025
- Bonas U, Schulte R, Fenselau S, Minsavage GV, Staskiewicz BJ, Stall RE (1991) Isolation of a gene cluster from *Xanthomonas campestris* pv. *vesicatoria* that determines pathogenicity and the hypersensitive response on pepper and tomato. *Mol Plant Microbe Interact* 4: 81-88
- Boucher CA, Van Gijsegem F, Barberis PA, Ayali M, Zischek C (1987) *Pseudomonas solanacearum* genes controlling both pathogenicity and hypersensitivity on tobacco are clustered. *J Bacteriol* 159: 5626-5632
- Boucher CA, Gough CL, Ayali M (1992) Molecular genetics of pathogenicity determinants of *Pseudomonas solanacearum* with special emphasis on *htr* genes. *Annu Rev Phytopathol* 30: 443-461
- Carpenter PG, Odel GW (1993) *Bacillus subtilis* *FlhA*, a flagellar protein related to a new family of signal transducing receptors. *Mol Microbiol* 7: 735-743
- Coplin DL, Frederick RD, Majerczak DR, Tuttle LD (1992) Characterization of a gene cluster that specifies pathogenicity in *Erwinia stewartii*. *Mol Plant Microbe Interact* 5: 81-88
- Cornelis G, Slutsky C, Lambert de Rouvroit C, Michiels T (1989) Homology between *Ynf*, the transcriptional activator of the *Yersinia* virulence regulon, and *AraC*, the *Escherichia coli* *arabinose* operon regulator. *J Bacteriol* 171: 254-262
- Fellay R, Nahne LG, Mindros MN, Frederick RD, Piel A, Panopoulos NJ (1991) Genes and signals controlling the *Pseudomonas syringae* pv. *phaseolicola*-plant interaction, in Hamecke H, Verma DPS (eds) *Advances in molecular genetics of plant-microbe interactions* vol 1, Kluwer Academic, Dordrecht, Netherlands, pp 45-52
- Fenselau S, Ballo I, Bonas U (1992) Determinants of pathogenicity in *Xanthomonas campestris* pv. *vesicatoria* are related to proteins involved in secretion in bacterial pathogens of animals. *Mol Plant-Microbe Interact* 4: 593-601
- Frederick RD, Majerczak DR, Coplin DL (1993) *Erwinia stewartii* *WspA*, a positive regulator of pathogenicity gene expression, is similar to *Pseudomonas syringae* pv. *phaseolicola* *HrpS*. *Mol Microbiol* 9: 477-485
- Galen JE, Curtiss III (1990) Expression of *Salmonella typhimurium* genes required for invasion is regulated by changes in DNA supercoiling. *Infect Immun* 58: 1879-1885
- Galán JE, Gincoldo C, Costas P (1992) Molecular and functional characterization of the *Salmonella* invasion gene *invA*: homology of *invA* to members of a new protein family. *J Bacteriol* 174: 4338-4349
- Gann S, Gough CL, Zischek C, Boucher CA (1989) The *htr* gene encodes a protein regulator of pathogenicity genes from *Pseudomonas solanacearum*. *Mol Microbiol* 6: 3005-3076
- Gann S, Gough CL, Ayali M, Zischek C, Van Gijsegem F, Barberis P, Boucher CA (1993) Involvement of *Pseudomonas solanacearum* *htr* genes in the secretion of a bacterial compound which induces a hypersensitive-like response on tobacco. In: Nestler EW, Verma DPS (eds) *Advances in molecular genetics of plant-microbe interactions*, vol 2, Kluwer Academic, Dordrecht, Netherlands, pp 259-268
- Gagnon JD, Yohrer J, Straley SC (1994) Genetic analysis of the low calcium response in *Yersinia* *pasteurii* *Mu* d1 *Yop* loci insertion mutants. *J Bacteriol* 160: 842-848
- Gough CL, Gann S, Zischek C, Boucher CA (1992) *htr* genes of *Pseudomonas solanacearum* are homologous to pathogenicity determinants of animal pathogenic bacteria and are conserved among plant pathogenic bacteria. *Mol Plant Microbe Interact* 5: 384-389
- Gough CL, Gann S, Lopes V, Boucher CA (1993) Homology between the *HrpO* protein of *Pseudomonas*

- solanacearum and bacterial proteins implicated in a signal peptide independent secretion mechanism. *Mol Gen Genet* 238: 378-382
- Grimm C, Panopoulos NJ (1989) The predicted product of a pathogenicity locus from *Pseudomonas syringae* pv. phaseolicola is homologous to a highly conserved domain of several prokaryotic regulatory proteins. *J Bacteriol* 171: 6031-6038
- Grissman EA, Ochman H (1993) Copraie gene clusters govern invasion of host epithelial cells by *Salmonella typhimurium* and *Shigella flexneri*. *EMBO J* 12: 3779-3787
- Haddix PL, Straley SC (1992) Structure and regulation of the *Yersinia pestis* *ysbCDEF* operon. *J Bacteriol* 174: 4820-4828
- Ilala TI, (1991) Genetic basis for virulence in *Shigella* species. *Microbiol Rev* 55: 208-224
- He SY, Huang HC, Collier A (1993) *Pseudomonas syringae* pv. *syringae* harpin, a protein that is secreted via the *Hrp* pathway and elicits the hypersensitive response in plants. *Cell* 73: 1255-1268
- Hou S, Hutcheson SW (1993) Nucleotide sequence and properties of the *hmrA* locus associated with the *Hrp* pathway. *Mol Plant Microbe Interact* 6: 553-564
- Huang HC, Schuurink R, Denny TP, Atkinson MM, Baker CJ, Yucel I, Hutcheson SW, Collier A (1988) Molecular cloning of a *Pseudomonas syringae* pv. *syringae* gene cluster that enables *Pseudomonas fluorescens* to elicit the hypersensitive response in tobacco plants. *J Bacteriol* 170: 4748-4758
- Huang Y, Xu P, Serpinin L (1990) A second cluster of genes that specify pathogenicity and host response in *Pseudomonas solanacearum*. *Mol Plant Microbe Interact* 3: 48-53
- Huang HC, Hutcheson SW, Collier A (1991) Characterization of the *hrp* cluster from *Pseudomonas syringae* pv. *syringae* 61 and *Trichobacterales* of genes encoding exported or membrane-spanning *Hrp* proteins. *Mol Plant Microbe Interact* 4: 469-478
- Huang HC, He SY, Bauer DW, Collier A (1992) The *Pseudomonas syringae* pv. *syringae* 61 *hrpA* product, an envelope protein required for elicitation of the hypersensitive response in plants. *J Bacteriol* 174: 6878-6885
- Huang HC, Xiao Y, Lin RH, Lu Y, Hutcheson SW, Collier A (1993) Characterization of the *Pseudomonas syringae* pv. *syringae* *hrpA* and *hrpI* genes: homology of *HrpI* to a superfamily of proteins associated with protein translocation. *Mol Plant Microbe Interact* 6: 515-520
- Huynh TV, Dahlbeck D, Staskiewicz BJ (1989) Bacterial blight of soybean: regulation of a pathogen gene determining host cultivar specificity. *Science* 245: 1374-1377
- Hwang I, Lin SM, Shaw PD (1992) Cloning and characterization of pathogenicity genes from *Xanthomonas campestris* pv. *glyceriae*. *J Bacteriol* 174: 1923-1931
- Jones RW, Bent AF, Kunkel BN, Bisgrove SR, Staskiewicz BJ (1993) Molecular analysis of virulence genes *avrPto2* and identification of a putative regulatory sequence common to all known *Pseudomonas syringae* virulence genes. *J Bacteriol* 175: 4859-4869
- Kamdar HV, Kamoun S, Kado CI (1993) Restoration of pathogenicity of avirulent *Xanthomonas oryzae* pv. *oryzae* and *X. campestris* pathogens by reciprocal complementation with the *hrpX* and *hrpXc* genes and identification of *HrpX* function by sequence analysis. *J Bacteriol* 175: 2017-2025
- Kamoun S, Kado CI (1990) A plant-inducible gene of *Xanthomonas campestris* pv. *campestris* encodes an extracellular component required for growth in the host and hypersensitivity on nonhosts. *J Bacteriol* 172: 5165-5172
- Kamoun S, Kamdar HV, Tolia E, Kado CI (1992) Incompatible interactions between clusters and *Xanthomonas campestris* involve a vascular hypersensitive response: role of the *hrpX* locus. *Mol Plant Microbe Interact* 5: 22-33
- Klement Z (1982) Hypersensitivity in: Mount MS, Lacey GH (eds) *Phytopathogenic prokaryotes*, vol 2. Academic Press, New York, pp 149-177
- Levy RJ, Beer SV (1992) Hybridization and functional complementation of the *hrp* gene cluster from *Erwinia amylovora* strain Ea321 with DNA of other bacteria. *Mol Plant Microbe Interact* 5: 412-419
- Lambert de Rouvroil C, Stuiters C, Cornelis GR (1992) Role of the transcriptional activator, *Vif*, and temperature in the expression of the *pYV* plasmid genes of *Yersinia enterocolica*. *Mol Microbiol* 6: 395-409
- Little RP, Mitsuhashi GV, Bonas U, Stoll RE (1994) Detection and identification of plant pathogenic strains of *Xanthomonas* based on amplification of DNA sequences related to the *hrp* genes of *Xanthomonas campestris* pv. *vesicatoria*. *Appl Environ Microbiol* 60: 1080-1077
- Linker PB, Peel RC, Panopoulos NJ (1986) Gene cluster of *Pseudomonas syringae* pv. *phaseolicola* controls pathogenicity on bean plants and hypersensitivity on nonhost plants. *J Bacteriol* 168: 5172-5177
- Linker PB, Panopoulos NJ, Staskiewicz BJ, Dahlbeck D (1988) Genes required for pathogenicity and hypersensitivity are conserved and intercalangulate among pathogens of *Pseudomonas syringae*. *Mol Gen Genet* 211: 499-506

- Linker WP, Lamb CJ, Dixon RA (1983) Microbial recognition and activation of plant defense systems. *Trends Microbiol* 1: 181-187
- Loudens L, Debussche L, Bon A, Lacroix JM, Bohn J-P (1983) Homology between a genetic locus (*hrpA*) involved in the osmoregulated biosynthesis of periplasmic glucans in *Escherichia coli* and a genetic locus (*hrpA*) controlling pathogenicity of *Pseudomonas syringae*. *Mol Microbiol* 10: 329-340
- Mekalanos JJ (1992) Environmental signals controlling expression of virulence determinants in bacteria. *J Bacteriol* 174: 1-7
- Meinhardt LW, Krishnan HB, Balwit PA, Puspoke SG (1993) Molecular cloning and characterization of a *sym* plasmid locus that regulates culture-specific nodulation of soybean by *Rhizobium loti* USDA257. *Mol Microbiol* 8: 17-29
- Michels T, Wain P, Bressan R, Ruysschaert JM, Cornelis G (1990) Secretion of Yop proteins by *Yersinia*. *Infect Immun* 58: 2840-2849
- Michels T, Vanooteghem J-C, Lambert de Rouvroil C, China B, Justin A, Baudry P, Cornelis GR (1991) Analysis of *vifC*, an operon involved in the secretion of Yop proteins by *Yersinia enterocolica*. *J Bacteriol* 173: 4594-5009
- Miller W, Mindrinos MS, Rahme LG, Frederick RD, Grimm C, Grossman R, Kyriakides X, Kottakis M, Panopoulos NJ (1993) *Pseudomonas syringae* pv. *phaseolicola*-plant interactions: host-pathogen signaling through cascade control of *hrp* gene expression. In: Nester EW, Verma DP (eds) *Advances in molecular genetics of plant-microbe interactions*, vol 2. Kluwer Academic, Dordrecht, pp 267-274
- Mukhopadhyay P, Williams J, Mills D (1988) Molecular analysis of a pathogenicity locus in *Pseudomonas syringae* pv. *syringae*. *J Bacteriol* 170: 5479-5488
- Mulholland V, Hinton JCD, Sidebottom J, Toth JK, Hyman LJ, Perombelon MCM, Reeves PJ, Salmon GPC (1993) A pleiotropic reduced virulence (*hrv1*) mutant of *Erwinia carotovora* subspecies *atopica* is defective in flagella assembly proteins that are conserved in plant and animal bacterial pathogens. *Mol Microbiol* 9: 343-356
- Nepold F, Anderson D, Mills D (1985) Cloning determinants of pathogenesis from *Pseudomonas syringae* pv. *syringae*. *Proc Natl Acad Sci USA* 82: 408-410
- Pano GV, Barve SS, Straley SC (1991) *LcdD*, a membrane-bound regulator of the *Yersinia pestis* low-calcium response. *J Bacteriol* 173: 7293-7303
- Rahme LG, Mindrinos MN, Panopoulos NJ (1991) Genetic and transcriptional organization of the *hrp* cluster of *Pseudomonas syringae* pv. *phaseolicola*. *J Bacteriol* 173: 575-586
- Rahme LG, Mindrinos MN, Panopoulos NJ (1992) Plant and environmental sensory signals control the expression of *hrp* genes in *Pseudomonas syringae* pv. *phaseolicola*. *J Bacteriol* 174: 3499-3507
- Ramkrishnan G, Zhu J-L, Newton A (1991) The cell cycle-regulated flagellar gene *flbF* of *Caulobacter crescentus* is homologous to a virulence locus (*hrpD*) of *Yersinia pestis*. *J Bacteriol* 173: 7283-7292
- Ruppelmann M, Gerstberg A, Wolf-Watz H (1992) A novel protein, *LcdD*, involved in the low-calcium response of *Yersinia pseudotuberculosis* shows extensive homology to YopI. *J Bacteriol* 174: 3355-3363
- Salmón JM, Staskiewicz BJ (1993) Molecular characterization and *hrp* dependence of the virulence gene *avrPto* from *Pseudomonas syringae* pv. *tomato*. *Mol Gen Genet* 239: 6-16
- Sanders LA, Van Way S, Mullin DA (1992) Characterization of the *Caulobacter crescentus* *flbF* promoter and identification of the inferred *FlbF* product as a homolog of the *LcdD* protein from a *Yersinia enterocolica* virulence plasmid. *J Bacteriol* 174: 857-868
- Sassie M, Gay NJ, Ebdale A, Runswick MJ, Walker JE (1981) The *alp* operon: nucleotide sequences of the genes for the a, b, and g subunits of *Escherichia coli* ATP synthase. *Nucleic Acids Res* 9: 5281-5296
- Sasakawa C, Komatsu K, Tobe T, Suzuki T, Yoshizawa M (1993) Eight genes in region 5 that form an operon are essential for invasion of epithelial cells by *Shigella flexneri*. *J Bacteriol* 175: 2334-2346
- Schulze R, Bonas U (1992a) Expression of the *Xanthomonas campestris* pv. *vesicatoria* *hrp* gene cluster, which determines pathogenicity and hypersensitivity on pepper and tomato, is plant inducible. *J Bacteriol* 174: 815-823
- Schulze R, Bonas U (1992b) A *Xanthomonas* pathogenicity locus is induced by sucrose and sulfur-containing amino acids. *Plant Cell* 4: 79-86
- Shim H, Kohn NI (1993) Characterization of the promoter of virulence gene D from *Pseudomonas syringae* pv. *tomato*. *J Bacteriol* 175: 5916-5924

Sinell JE, Minsavage GV (1990) The use of hrp genes to identify opportunistic xanthomonads. In: Klement Z (ed) plant pathogenic bacteria, proceedings of the 7th international conference of plant pathogenic bacteria, Budapest, Hungary, 1989. Akadémiai Kiadó, Budapest, pp 369-374

Steinberger EM, Beer SV (1998) Creation and complementation of pathogenicity mutants of *Erwinia amylovora*. Mol Plant Microbe Interact 1: 135-144

Straley SC, Plano GV, Strzpek E, Haddix PL, Fields KA (1993) Regulation by  $\text{Ca}^{2+}$  in the *Versinia lwo-*  $\text{Ca}^{2+}$  response. Mol Microbiol B: 1005-1010

Van Giesegem F, Genin S, Boucher C (1993) Conservation of secretion pathways for pathogenicity determinants of plant and animal bacteria. Trends Microbiol 1: 175-180

Venturiates MW, Brynes JM, Oaks EV (1992) Surface presentation of Shigella flexneri invasion plasmid antigens requires the products of the spa locus. J Bacteriol 174: 1990-2001

Vogler AP, Honma M, Iikura VM, Macrae RM (1991) Salmonella typhimurium mutants defective in flagellar filament regrowth and sequence similarity of Fil to FOF1, vacuolar, and archaeobacterial ATPase subunits. J Bacteriol 173: 3564-3572

Walters K, Marcoll A, Hitchlin E, Mansfield J (1990) Gene for pathogenicity and ability to cause hypersensitive reaction cloned from *Erwinia amylovora*. Physiol Mol Plant Pathol 36: 509-521

Weney VR, Kingsley MT, Gabriel DW (1991) Xanthomonas campestris pv. vitansucans genes determining host-specific virulence and general virulence on cereals identified by Tn5-quasi insertion mutagenesis. Mol Plant Microbe Interactions 4: 673-677

Weil ZM, Leiby RJ, Zumoff CH, Bauer DV, He SY, Collier A, Beer SV (1992a) Harpin, elicitor of the hypersensitive response produced by the plant pathogen *Erwinia amylovora*. Science 257: 85-89

Weil ZM, Sneath BJ, Beer SV (1992b) Expression of *Erwinia amylovora* hrp genes in response to environmental stimuli. J Bacteriol 174: 1875-1882

Weil ZM and Beer SV (1993) Hrp1 of *Erwinia amylovora* functions in secretion of harpin and is a member of a new protein family. J Bacteriol 175: 7958-7967

Willits DK, Rich JJ, Harbak EM (1991) hrp genes of phytopathogenic bacteria. Mol Plant Microbe Interact 4: 132-138

Wuans SC (1992) Two way chemical signaling in Agrobacterium-plant interactions. Microbiol Rev 58: 12-31

Xiao YX, Lu Y, Heu SG, Hutcheson SW (1992) Organization and environmental regulation of the *Pseudomonas syringae* pv. *syringae* 81 hrp cluster. J Bacteriol 174: 1734-1741

## 1 Action at the "Pathogenic Cusp"

Max-Deibüch-Laboratory, Carl-von-Linné-Weg 10, 50829 Köln, Germany